

SCIENCE

Data Booklet ***Revised September 2009***

Alberta
Education

Table of Contents

<i>Page</i>	
<i>1</i>	General Formulas and Data
	Units and Prefixes
<i>2</i>	Kinematics and Dynamics Formulas
	Gravitational and Electric Fields
	Astronomy Data
<i>3</i>	Electricity Formulas
	Wave Formulas
<i>4</i>	Electrochemistry
	Geological Time-Line
	Thermodynamics
<i>6</i>	Periodic Chart of the Elements and Ions
<i>8</i>	Nuclear Chemistry
<i>9</i>	Organic Chemistry
<i>10</i>	Solutions
<i>11</i>	Acids and Bases
<i>13</i>	Genetics
<i>14</i>	Scoring Descriptions for Standards Setting

Cover design interpretation of DNA in the presence of electromagnetic energy by Nathan A. Smith of Alberta Education.

Copyright 2009, the Crown in Right of Alberta, as represented by the Minister of Education, Alberta Education, Learner Assessment, 44 Capital Boulevard, 10044 108 Street NW, Edmonton, Alberta T5J 5E6, and its licensors. All rights reserved.

Special permission is granted to **Alberta educators only** to reproduce, for educational purposes and on a non-profit basis, this document or any of its parts.

General Formulas and Data

Formulas and Data

$$\text{slope} = \frac{\text{rise}}{\text{run}} = \frac{\Delta y}{\Delta x} = \frac{y_2 - y_1}{x_2 - x_1}$$

$$\text{percent difference from theoretical value} = \frac{\text{experimental value} - \text{theoretical value}}{\text{theoretical value}} \times 100\%$$

$$\text{percent efficiency} = \left(\frac{\text{output}}{\text{input}} \right) \times 100\%$$

$$\text{magnification} = \left(\text{power of ocular lens} \right) \times \left(\text{power of objective lens} \right)$$

**Distilled Water at
Room Temperature (25°C)
and Standard Pressure
(101.325 kPa)**

Volume	Mass	Density
1.0 mL or 1.0 cm ³	1.0 g	1.0 g/cm ³
1.0 L or 1.0 dm ³	1.0 kg	1.0 kg/dm ³

Units and Prefixes

Prefix	Symbol	Factor by Which Base Unit Is Multiplied	
tera	T	1 000 000 000 000	= 10 ¹²
giga	G	1 000 000 000	= 10 ⁹
mega	M	1 000 000	= 10 ⁶
kilo	k	1 000	= 10 ³
hecto	h	100	= 10 ²
deca	da	10	= 10 ¹
Common Base Units*		1	= 10 ⁰
deci	d	0.1	= 10 ⁻¹
centi	c	0.01	= 10 ⁻²
milli	m	0.001	= 10 ⁻³
micro	μ	0.000 001	= 10 ⁻⁶
nano	n	0.000 000 001	= 10 ⁻⁹
pico	p	0.000 000 000 001	= 10 ⁻¹²

*metre (m), gram (g), litre (L), mole (mol)

Some Non-SI Units Used with SI

Quantity	Unit Name	Symbol	Definition
Time	minute	min	1 min = 60 s
	hour	h	1 h = 3 600 s
	day	d	1 d = 86 400 s
	year (annum)	a	1 a = 31 557 600 s
Area	hectare	ha	1 ha = 1 hm ² = 10 000 m ²
Volume	litre	L	1 L = 1 000 cm ³
Mass	metric ton or tonne	t	1 t = 1 000 kg = 1 Mg
Pressure	standard atmosphere	atm	1 atm = 101.325 kPa

Kinematics and Dynamics Formulas

$$v = \frac{\Delta d}{\Delta t}$$

$$\vec{v} = \frac{\Delta \vec{d}}{\Delta t}$$

$$\vec{a} = \frac{\Delta \vec{v}}{\Delta t} = \frac{\vec{v}_f - \vec{v}_i}{\Delta t}$$

$$\vec{F}_{\text{net}} = m\vec{a}$$

$$\vec{F}_{\text{net}} = \vec{F}_a + \vec{F}_f$$

$$W = F\Delta d$$

$$P = \frac{W}{t}$$

$$\Delta \vec{d} = \vec{v}_i \Delta t + \frac{1}{2} \vec{a} (\Delta t)^2$$

$$\Delta \vec{d} = \left(\frac{\vec{v}_i + \vec{v}_f}{2} \right) \Delta t$$

$$\vec{p} = m\vec{v}$$

$$\Delta \vec{p} = \vec{F} \Delta t, \Delta \vec{p} = \vec{p}_f - \vec{p}_i$$

$$\vec{F} = \frac{m(\vec{v}_f - \vec{v}_i)}{\Delta t}$$

$$E_p = mgh$$

$$E_k = \frac{1}{2}mv^2$$

v = average speed (m/s)

$\vec{v}$ = average velocity (m/s)

d = distance (m)

$\vec{d}$ = displacement (m)

t = time elapsed (s)

$\vec{a}$ = acceleration (m/s²)

$\vec{F}$ = force (kg·m/s² or N)

$\vec{F}_{\text{net}}$ = net force (N)

$\vec{F}_a$ = applied force (N)

$\vec{F}_f$ = force of friction (N)

F = magnitude of a force (N)

m = mass (kg)

W = work (N·m or J)

P = power (J/s or W)

Δ = change in

$\vec{F} \Delta t$ = impulse

$\vec{p}$ = momentum (kg·m/s)

E_p = gravitational potential energy (J)

g = magnitude of acceleration due to gravity (m/s²)

E_k = kinetic energy (J)

Collisions

Hit and rebound:

$$m_1 \vec{v}_1 + m_2 \vec{v}_2 = m_1 \vec{v}'_1 + m_2 \vec{v}'_2$$

Hit and stick:

$$m_1 \vec{v}_1 + m_2 \vec{v}_2 = (m_1 + m_2) \vec{v}'_{1 \text{ and } 2}$$

Explosion:

$$(m_1 + m_2) \vec{v}_{1 \text{ and } 2} = m_1 \vec{v}'_1 + m_2 \vec{v}'_2$$

Gravitational and Electric Fields

$$\vec{F}_g = m\vec{g}$$

$$g = \frac{Gm}{r^2}$$

$$|\vec{E}| = \frac{kq}{r^2}$$

$\vec{F}_g$ = force due to gravity (N)

m = mass (kg)

G = gravitational constant = $6.67 \times 10^{-11} \text{ N} \cdot \text{m}^2/\text{kg}^2$

r = radius or centre-to-centre distance (m)

g = magnitude of gravitational field strength (N/kg)

k = Coulomb's law constant = $8.99 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2$

q = electrostatic charge in coulombs (C)

$|\vec{E}|$ = electric field strength (N/C)

Astronomy Data

Mass of Earth = $5.98 \times 10^{24} \text{ kg}$

Radius of Earth = $6.37 \times 10^6 \text{ m}$

Mass of sun = $1.99 \times 10^{30} \text{ kg}$

1 light-year = $9.47 \times 10^{15} \text{ m}$

1 AU (astronomical unit) = $1.50 \times 10^{11} \text{ m}$

Average acceleration due

to gravity on surface of Earth = 9.81 m/s^2

Average gravitational field

strength on surface of Earth = 9.81 N/kg

Electricity Formulas

$$P = IV, \quad P = I^2 R$$

$$V = IR$$

$$E = Pt$$

For resistances connected in series

$$R_T = R_1 + R_2 + R_3 + \dots R_n$$

For resistances connected in parallel

$$\frac{1}{R_T} = \frac{1}{R_1} + \frac{1}{R_2} + \frac{1}{R_3} + \dots \frac{1}{R_n}$$

Ideal Transformers

$$\frac{N_p}{N_s} = \frac{V_p}{V_s}, \quad \frac{N_p}{N_s} = \frac{I_s}{I_p}, \quad \frac{V_p}{V_s} = \frac{I_s}{I_p}$$

R = resistance (Ω)

P = power (W)

I = current (A)

V = voltage (V)

E = energy (J)

t = time elapsed (s)

N = number of turns

p = primary

s = secondary

Related value: 1.00 kilowatt hour = 1.00 kW·h = 3.60×10^6 J

Wave Formulas

$$v = f\lambda$$

$$c = f\lambda$$


v = speed of wave (m/s)

c = speed of electromagnetic radiation in air or vacuum (3.00×10^8 m/s)

f = frequency (Hz or 1/s)

λ = wavelength (m)

Electromagnetic Spectrum



Electrochemistry

**Activity Series for 1.0 mol/L Solution
at 25 °C and 101.325 kPa**

	Reduction Half-Reaction				
Increasing strength of reactant as an oxidizing agent ↑	Au ³⁺ (aq)	+	3 e ⁻	→	Au(s)
	Hg ²⁺ (aq)	+	2 e ⁻	→	Hg(l)
	Ag ⁺ (aq)	+	e ⁻	→	Ag(s)
	Cu ²⁺ (aq)	+	2 e ⁻	→	Cu(s)
	2 H ⁺ (aq)	+	2 e ⁻	→	H ₂ (g)
	Pb ²⁺ (aq)	+	2 e ⁻	→	Pb(s)
	Sn ²⁺ (aq)	+	2 e ⁻	→	Sn(s)
	Ni ²⁺ (aq)	+	2 e ⁻	→	Ni(s)
	Cd ²⁺ (aq)	+	2 e ⁻	→	Cd(s)
	Fe ²⁺ (aq)	+	2 e ⁻	→	Fe(s)
	Zn ²⁺ (aq)	+	2 e ⁻	→	Zn(s)
	Cr ²⁺ (aq)	+	2 e ⁻	→	Cr(s)
	Al ³⁺ (aq)	+	3 e ⁻	→	Al(s)
	Mg ²⁺ (aq)	+	2 e ⁻	→	Mg(s)
	Na ⁺ (aq)	+	e ⁻	→	Na(s)
	Ca ²⁺ (aq)	+	2 e ⁻	→	Ca(s)
	Li ⁺ (aq)	+	e ⁻	→	Li(s)
					Increasing strength of reactant as a reducing agent ↓

Thermodynamics

Heat Capacities of Selected Substances at 25 °C

Compound		Specific Heat Capacity (J/g·°C) or (kJ/kg·°C)
water	H ₂ O(l)	4.19
ice (at 0 °C)	H ₂ O(s)	2.10
water vapour (at 100 °C)	H ₂ O(g)	2.08
methanol	CH ₃ OH(l)	2.53
ethanol	C ₂ H ₅ OH(l)	2.44
hexane	C ₆ H ₁₄ (l)	2.27
toluene	C ₇ H ₈ (l)	1.71
air	mixture of N ₂ (g), O ₂ (g), CO ₂ (g), and trace gases	1.01

Thermodynamic Properties of Selected Compounds

Compound		Melting Point (°C)	Boiling Point (°C)	Heat of Fusion (kJ/mol)	Heat of Vaporization (kJ/mol)
water	H ₂ O(l)	0.00	100.00	6.01	40.66
hexane	C ₆ H ₁₄ (l)	-95.35	68.73	13.08	28.85
ethanol	C ₂ H ₅ OH(l)	-114.14	78.29	4.93	38.56
methanol	CH ₃ OH(l)	-97.53	64.6	3.22	35.21
toluene	C ₇ H ₈ (l)	-94.95	110.63	6.64	33.18

Geological Time-Line

Millions of Years Ago	Era	Period	Epoch
1.7	Cenozoic	Quaternary	Holocene Pleistocene
	Mesozoic		Tertiary
65			
			Cretaceous
140			
	Paleozoic		Jurassic
210			
			Triassic
250			
290			Permian
			Carboniferous
360			
			Devonian
410			
440			Silurian
	Precambrian		Ordovician
500			
			Cambrian
590			
4 600			

Standard Heats of Formation of Selected Compounds at 25°C

Compound	Formula	$\Delta_f H^\circ$ (kJ/mol)
ammonia	$\text{NH}_3(\text{g})$	-45.9
benzene	$\text{C}_6\text{H}_6(\text{l})$	+49.1
butane	$\text{C}_4\text{H}_{10}(\text{g})$	-125.7
calcium carbonate	$\text{CaCO}_3(\text{s})$	-1 207.6
calcium hydroxide	$\text{Ca}(\text{OH})_2(\text{s})$	-985.2
carbon dioxide	$\text{CO}_2(\text{g})$	-393.5
carbon monoxide	$\text{CO}(\text{g})$	-110.5
ethane	$\text{C}_2\text{H}_6(\text{g})$	-84.0
ethanoic acid (acetic acid)	$\text{CH}_3\text{COOH}(\text{l})$	-484.3
ethanol	$\text{C}_2\text{H}_5\text{OH}(\text{l})$	-277.6
ethene (ethylene)	$\text{C}_2\text{H}_4(\text{g})$	+52.4
ethyne (acetylene)	$\text{C}_2\text{H}_2(\text{g})$	+227.4
glucose	$\text{C}_6\text{H}_{12}\text{O}_6(\text{s})$	-1 273.3
hydrogen sulfide	$\text{H}_2\text{S}(\text{g})$	-20.6
methane	$\text{CH}_4(\text{g})$	-74.6
methanol	$\text{CH}_3\text{OH}(\text{l})$	-239.2
nitrogen dioxide	$\text{NO}_2(\text{g})$	+33.2
nitrogen monoxide	$\text{NO}(\text{g})$	+91.3
octane	$\text{C}_8\text{H}_{18}(\text{l})$	-250.1
pentane	$\text{C}_5\text{H}_{12}(\text{l})$	-173.5
propane	$\text{C}_3\text{H}_8(\text{g})$	-103.8
sucrose	$\text{C}_{12}\text{H}_{22}\text{O}_{11}(\text{s})$	-2 226.1
sulfur dioxide	$\text{SO}_2(\text{g})$	-296.8
sulfur trioxide	$\text{SO}_3(\text{g})$	-395.7
water (liquid)	$\text{H}_2\text{O}(\text{l})$	-285.8
water (gas)	$\text{H}_2\text{O}(\text{g})$	-241.8

Note: Elements are given a value of zero.
 Negative sign (-) denotes exothermic change.
 Positive sign (+) denotes endothermic change.

Energy Formulas

$$Q = mc\Delta t$$

$$\Delta_{\text{fus}} H = \frac{Q}{n}$$

$$\Delta_{\text{vap}} H = \frac{Q}{n}$$

$$\Delta_r H = \sum n \Delta_f H^\circ \text{ products} - \sum n \Delta_f H^\circ \text{ reactants}$$

Q = quantity of heat energy (J or kJ)

m = mass (g or kg)

$\Delta_{\text{fus}} H$ = heat of fusion (kJ/mol)

$\Delta_{\text{vap}} H$ = heat of vaporization (kJ/mol)

c = specific heat capacity (J/g·°C or kJ/kg·°C)

Δt = change in temperature (°C)

n = amount in moles (mol)

$\Delta_r H$ = energy change of reaction (kJ)

Σ = the sum of

$\Delta_f H^\circ$ = standard molar heat (enthalpy) of formation (kJ/mol)

Periodic Chart of the Elements and Ions

1	2	3	4	5	6	7	8	9
---	---	---	---	---	---	---	---	---

1 H hydrogen 1.01 H ⁺ hydrogen	
3 Li lithium 6.94 Li ⁺ lithium	4 Be beryllium 9.01 Be ²⁺ beryllium
11 Na sodium 22.99 Na ⁺ sodium	12 Mg magnesium 24.31 Mg ²⁺ magnesium

Note: The legend at the right denotes the physical state of the elements at 101.325 kPa and 298.15 K (25°C).

Legend for the Elements

Solid	Liquid	Gas	Seldom forms ions
-------	--------	-----	-------------------

Table of Polyatomic Ions

Polyatomic ions									
acetate	CH ₃ COO ⁻	hydrogen carbonate	HCO ₃ ⁻	cyanide	CN ⁻	phosphate	PO ₄ ³⁻	sulfite	SO ₃ ²⁻
ammonium	NH ₄ ⁺	chlorate	ClO ₃ ⁻	hydroxide	OH ⁻	hydrogen phosphate	HPO ₄ ²⁻	hydrogen sulfide	HS ⁻
benzoate	C ₆ H ₅ COO ⁻	hypochlorite	ClO ⁻	nitrate	NO ₃ ⁻	dihydrogen phosphate	H ₂ PO ₄ ⁻	hydrogen sulfate	HSO ₄ ⁻
borate	BO ₃ ³⁻	chromate	CrO ₄ ²⁻	nitrite	NO ₂ ⁻	silicate	SiO ₃ ²⁻	hydrogen sulfite	HSO ₃ ⁻
carbonate	CO ₃ ²⁻	dichromate	Cr ₂ O ₇ ²⁻	permanganate	MnO ₄ ⁻	sulfate	SO ₄ ²⁻		

19 K potassium 39.10 K ⁺ potassium	20 Ca calcium 40.08 Ca ²⁺ calcium	21 Sc scandium 44.96 Sc ³⁺ scandium	22 Ti titanium 47.87 Ti ⁴⁺ titanium(IV) Ti ³⁺ titanium(III)	23 V vanadium 50.94 V ⁵⁺ vanadium(V) V ⁴⁺ vanadium(IV)	24 Cr chromium 52.00 Cr ³⁺ chromium(III) Cr ²⁺ chromium(II)	25 Mn manganese 54.94 Mn ²⁺ manganese(II) Mn ⁴⁺ manganese(IV)	26 Fe iron 55.85 Fe ³⁺ iron(III) Fe ²⁺ iron(II)	27 Co cobalt 58.93 Co ²⁺ cobalt(II) Co ³⁺ cobalt(III)
37 Rb rubidium 85.47 Rb ⁺ rubidium	38 Sr strontium 87.62 Sr ²⁺ strontium	39 Y yttrium 88.91 Y ³⁺ yttrium	40 Zr zirconium 91.22 Zr ⁴⁺ zirconium	41 Nb niobium 92.91 Nb ⁵⁺ niobium(V) Nb ³⁺ niobium(III)	42 Mo molybdenum 95.94 Mo ⁶⁺ molybdenum	43 Tc technetium (98) Tc ⁷⁺ technetium	44 Ru ruthenium 101.07 Ru ³⁺ ruthenium(III)	45 Rh rhodium 102.91 Rh ³⁺ rhodium
55 Cs cesium 132.91 Cs ⁺ cesium	56 Ba barium 137.33 Ba ²⁺ barium	57 La lanthanum 138.91 La ³⁺ lanthanum	72 Hf hafnium 178.49 Hf ⁴⁺ hafnium	73 Ta tantalum 180.95 Ta ⁵⁺ tantalum	74 W tungsten 183.84 W ⁶⁺ tungsten	75 Re rhenium 186.21 Re ⁷⁺ rhenium	76 Os osmium 190.23 Os ⁴⁺ osmium	77 Ir iridium 192.22 Ir ⁴⁺ iridium
87 Fr francium (223) Fr ⁺ francium	88 Ra radium (226) Ra ²⁺ radium	89 Ac actinium (227) Ac ³⁺ actinium	104 Rf rutherfordium (261)	105 Db dubnium (262)	106 Sg seaborgium (266)	107 Bh bohrium (264)	108 Hs hassium (277)	109 Mt meitnerium (268)

Lanthanide and Actinide Series Begins

Key	
Atomic number →	91 Pa
Name of the element →	protactinium
Atomic mass →	231.04
	Pa ⁵⁺
	protactinium(V)
	Pa ⁴⁺
	protactinium(IV)

Based on ¹²₆C

Most stable or common ion is listed above dotted line.
() indicates mass of the most stable isotope.

References

Lide, D.R. 2005. *CRC Handbook of Chemistry and Physics*. 86th ed. Boca Raton: CRC Press

58 Ce cerium 140.12 Ce ³⁺ cerium	59 Pr praseodymium 140.91 Pr ³⁺ praseodymium	60 Nd neodymium 144.24 Nd ³⁺ neodymium	61 Pm promethium (145) Pm ³⁺ promethium	62 Sm samarium 150.36 Sm ³⁺ samarium(III) Sm ²⁺ samarium(II)
90 Th thorium 232.04 Th ⁴⁺ thorium	91 Pa protactinium 231.04 Pa ⁵⁺ protactinium(V) Pa ⁴⁺ protactinium(IV)	92 U uranium 238.03 U ⁶⁺ uranium(VI) U ⁴⁺ uranium(IV)	93 Np neptunium (237) Np ⁵⁺ neptunium	94 Pu plutonium (244) Pu ⁴⁺ plutonium(IV) Pu ⁶⁺ plutonium(VI)

10	11	12	13	14	15	16	17	18
----	----	----	----	----	----	----	----	----

Polyatomic Elements

Elements			
astatine	At ₂	iodine	I ₂
bromine	Br ₂	nitrogen	N ₂
chlorine	Cl ₂	oxygen	O ₂
fluorine	F ₂	phosphorus	P ₄
hydrogen	H ₂	sulfur	S ₈

Polyatomic Elements

Elements			
astatine	At ₂	iodine	I ₂
bromine	Br ₂	nitrogen	N ₂
chlorine	Cl ₂	oxygen	O ₂
fluorine	F ₂	phosphorus	P ₄
hydrogen	H ₂	sulfur	S ₈

								1 hydrogen 1.01 H ⁻ hydride	2 helium 4.00 He helium				
								5 boron 10.81 B boron	6 carbon 12.01 C carbon	7 nitrogen 14.01 N ³⁻ nitride	8 oxygen 16.00 O ²⁻ oxide	9 fluorine 19.00 F ⁻ fluoride	10 neon 20.18 Ne neon
								13 aluminium 26.98 Al ³⁺ aluminium	14 silicon 28.09 Si silicon	15 phosphorus 30.97 P ³⁻ phosphide	16 sulfur 32.07 S ²⁻ sulfide	17 chlorine 35.45 Cl ⁻ chloride	18 argon 39.95 Ar argon
28 nickel 58.69 Ni ²⁺ nickel(II) Ni ³⁺ nickel(III)	29 copper 63.55 Cu ²⁺ copper(II) Cu ⁺ copper(I)	30 zinc 65.41 Zn ²⁺ zinc	31 gallium 69.72 Ga ³⁺ gallium	32 germanium 72.64 Ge ⁴⁺ germanium	33 arsenic 74.92 As ³⁻ arsenide	34 selenium 78.96 Se ²⁻ selenide	35 bromine 79.90 Br ⁻ bromide	36 krypton 83.80 Kr krypton					
46 palladium 106.42 Pd ²⁺ palladium(II) Pd ³⁺ palladium(III)	47 silver 107.87 Ag ⁺ silver	48 cadmium 112.41 Cd ²⁺ cadmium	49 indium 114.82 In ³⁺ indium	50 tin 118.71 Sn ⁴⁺ tin(IV) Sn ²⁺ tin(II)	51 antimony 121.76 Sb ³⁺ antimony(III) Sb ⁵⁺ antimony(V)	52 tellurium 127.60 Te ²⁻ telluride	53 iodine 126.90 I ⁻ iodide	54 xenon 131.29 Xe xenon					
78 platinum 195.08 Pt ⁴⁺ platinum(IV) Pt ²⁺ platinum(II)	79 gold 196.97 Au ³⁺ gold(III) Au ⁺ gold(I)	80 mercury 200.59 Hg ²⁺ mercury(II) Hg ⁺ mercury(I)	81 thallium 204.38 Tl ⁺ thallium(I) Tl ³⁺ thallium(III)	82 lead 207.2* Pb ²⁺ lead(II) Pb ⁴⁺ lead(IV)	83 bismuth 208.98 Bi ³⁺ bismuth(III) Bi ⁵⁺ bismuth(V)	84 polonium (209) Po ²⁺ polonium(II) Po ⁴⁺ polonium(IV)	85 astatine (210) At ⁻ astatide	86 radon (222) Rn radon					
110 darmstadtium (271)	111 roentgenium (272)												

* The isotopic mix of naturally occurring lead is more variable than that of other elements, preventing precision to greater than tenths of a gram per mole.

63 europium 151.96 Eu ³⁺ europium(III) Eu ²⁺ europium(II)	64 gadolinium 157.25 Gd ³⁺ gadolinium	65 terbium 158.93 Tb ³⁺ terbium	66 dysprosium 162.50 Dy ³⁺ dysprosium	67 holmium 164.93 Ho ³⁺ holmium	68 erbium 167.26 Er ³⁺ erbium	69 thulium 168.93 Tm ³⁺ thulium	70 ytterbium 173.04 Yb ³⁺ ytterbium(III) Yb ²⁺ ytterbium(II)	71 lutetium 174.97 Lu ³⁺ lutetium
95 americium (243) Am ³⁺ americium(III) Am ⁴⁺ americium(IV)	96 curium (247) Cm ³⁺ curium	97 berkelium (247) Bk ³⁺ berkelium(III) Bk ⁴⁺ berkelium(IV)	98 californium (251) Cf ³⁺ californium	99 einsteinium (252) Es ³⁺ einsteinium	100 fermium (257) Fm ³⁺ fermium	101 mendelevium (258) Md ²⁺ mendelevium(II) Md ³⁺ mendelevium(III)	102 nobelium (259) No ²⁺ nobelium(II) No ³⁺ nobelium(III)	103 lawrencium (262) Lr ³⁺ lawrencium

Nuclear Chemistry

Masses of Subatomic Particles and Radiation

Subatomic Particle or Radiation	Mass (10^{-3} kg/mol)	Subatomic Particle or Radiation	Mass (10^{-3} kg/mol)
alpha particle (helium nucleus) ${}^4_2\text{He}$ or α	4.001 51	positron ${}^0_{+1}\text{e}$	0.000 549
beta particle (electron) ${}^0_{-1}\text{e}$ or β	0.000 549	gamma radiation ${}^0_0\gamma$	—
		neutron ${}^1_0\text{n}$	1.008 66
		proton ${}^1_1\text{p}$	1.007 28

Masses of Selected Nuclides

Nuclide	Mass (10^{-3} kg/mol)	Nuclide	Mass (10^{-3} kg/mol)
barium-141 ${}^{141}_{56}\text{Ba}$	140.914 41	nitrogen-15 ${}^{15}_7\text{N}$	15.000 11
beryllium-7 ${}^7_4\text{Be}$	7.016 93	oxygen-15 ${}^{15}_8\text{O}$	15.003 07
beryllium-8 ${}^8_4\text{Be}$	8.005 31	oxygen-16 ${}^{16}_8\text{O}$	15.994 91
boron-8 ${}^8_5\text{B}$	8.024 61	oxygen-18 ${}^{18}_8\text{O}$	17.999 16
carbon-14 ${}^{14}_6\text{C}$	14.003 24	phosphorus-31 ${}^{31}_{15}\text{P}$	30.973 76
cesium-144 ${}^{144}_{55}\text{Cs}$	143.932 02	plutonium-239 ${}^{239}_{94}\text{Pu}$	239.052 16
fluorine-17 ${}^{17}_9\text{F}$	17.002 10	polonium-210 ${}^{210}_{84}\text{Po}$	209.982 86
helium-3 ${}^3_2\text{He}$	3.016 03	polonium-218 ${}^{218}_{84}\text{Po}$	218.008 97
hydrogen-1 ${}^1_1\text{H}$	1.007 83	potassium-40 ${}^{40}_{19}\text{K}$	39.964 00
hydrogen-2 (deuterium) ${}^2_1\text{H}$	2.014 10	radium-226 ${}^{226}_{88}\text{Ra}$	226.025 40
hydrogen-3 (tritium) ${}^3_1\text{H}$	3.016 03	radon-222 ${}^{222}_{86}\text{Rn}$	222.017 57
krypton-92 ${}^{92}_{36}\text{Kr}$	91.926 11	rubidium-90 ${}^{90}_{37}\text{Rb}$	89.914 81
lanthanum-146 ${}^{146}_{57}\text{La}$	145.925 8	ruthenium-107 ${}^{107}_{44}\text{Ru}$	106.909 9
lead-206 ${}^{206}_{82}\text{Pb}$	205.974 5	strontium-95 ${}^{95}_{38}\text{Sr}$	94.919 31
lead-208 ${}^{208}_{82}\text{Pb}$	207.976 64	sulfur-31 ${}^{31}_{16}\text{S}$	30.979 56
neon-20 ${}^{20}_{10}\text{Ne}$	19.992 44	thorium-230 ${}^{230}_{90}\text{Th}$	230.033 13
nitrogen-13 ${}^{13}_7\text{N}$	13.005 74	uranium-235 ${}^{235}_{92}\text{U}$	235.043 92
nitrogen-14 ${}^{14}_7\text{N}$	14.003 07		

Elements for Radioactive Dating

Radioisotope (Parent Nuclide)	Final Decay Nuclide	Approximate Half-Life (annum—a)
carbon-14 ${}^{14}_6\text{C}$	nitrogen-14 ${}^{14}_7\text{N}$	5.73×10^3
potassium-40 ${}^{40}_{19}\text{K}$	argon-40 ${}^{40}_{18}\text{Ar}$	1.26×10^9
rubidium-87 ${}^{87}_{37}\text{Rb}$	strontium-87 ${}^{87}_{38}\text{Sr}$	4.88×10^{10}
uranium-235 ${}^{235}_{92}\text{U}$	lead-207 ${}^{207}_{82}\text{Pb}$	7.04×10^8
uranium-238 ${}^{238}_{92}\text{U}$	lead-206 ${}^{206}_{82}\text{Pb}$	4.47×10^9


Energy Change Formula

$$\Delta E = \Delta mc^2$$

ΔE = change in energy (J)

Δm = mass converted to energy (kg)

c = speed of EMR (3.00×10^8 m/s)



Organic Chemistry

Homologous Series of Alkanes at 25°C and 101.325 kPa

Name*	Formula	Name*	Formula
methane	CH ₄ (g)	hexane	C ₆ H ₁₄ (l)
ethane	C ₂ H ₆ (g)	heptane	C ₇ H ₁₆ (l)
propane	C ₃ H ₈ (g)	octane	C ₈ H ₁₈ (l)
butane	C ₄ H ₁₀ (g)	nonane	C ₉ H ₂₀ (l)
pentane	C ₅ H ₁₂ (l)	decane	C ₁₀ H ₂₂ (l)

*Note: Italics indicate organic nomenclature prefixes.

Prefixes for Molecular Compounds

1 = mono	6 = hexa-
2 = di-	7 = hepta-
3 = tri-	8 = octa-
4 = tetra-	9 = ennea- (nona-)
5 = penta-	10 = deca-

Types of Reactions

Formation (Synthesis)

element + element → compound

Decomposition

compound → element + element

Single Replacement

compound + element → new compound + new element

Double Replacement

compound + compound → new compound + new compound

Complete Hydrocarbon Combustion

hydrocarbon + oxygen → carbon dioxide + water

Addition

alkene or alkyne + hydrogen → alkane

alkene or alkyne + halogen → halogenated hydrocarbon

Cracking

large hydrocarbon → small hydrocarbon

Polymerization

monomer + monomer → polymer

Esterification

alcohol + carboxylic acid → ester + water

General Formulas and Names of Some Organic Compounds

General Formula	Classification	Example Formula	Example Name
C _n H _(2n+2)	alkane	$ \begin{array}{c} \text{H} \quad \text{H} \\ \quad \\ \text{H}-\text{C}-\text{C}-\text{H} \\ \quad \\ \text{H} \quad \text{H} \end{array} $	ethane
C _n H _(2n)	alkene	$ \begin{array}{c} \text{H} \quad \text{H} \\ \diagdown \quad \diagup \\ \text{C} = \text{C} \\ \diagup \quad \diagdown \\ \text{H} \quad \text{H} \end{array} $	ethene
C _n H _(2n-2)	alkyne	H - C ≡ C - H	ethyne
R - O - H	alcohol	$ \begin{array}{c} \text{H} \quad \text{H} \\ \quad \\ \text{H}-\text{C}-\text{C}-\text{O}-\text{H} \\ \quad \\ \text{H} \quad \text{H} \end{array} $	ethanol
$ \begin{array}{c} \text{O} \\ \\ \text{R}-\text{C} \\ \\ \text{O}-\text{H} \end{array} $	carboxylic acid	$ \begin{array}{c} \text{H} \quad \text{O} \\ \quad \\ \text{H}-\text{C}-\text{C} \\ \quad \\ \text{H} \quad \text{O}-\text{H} \end{array} $	ethanoic acid
$ \begin{array}{c} \text{O} \\ \\ \text{R}-\text{C} \\ \\ \text{O}-\text{R}' \end{array} $	ester	$ \begin{array}{c} \text{H} \quad \text{O} \quad \text{H} \\ \quad \quad \\ \text{H}-\text{C}-\text{C}-\text{O}-\text{C}-\text{H} \\ \quad \quad \\ \text{H} \quad \quad \text{H} \end{array} $	methyl ethanoate
R - Q	halogenated hydrocarbon	$ \begin{array}{c} \text{H} \quad \text{H} \\ \quad \\ \text{H}-\text{C}-\text{C}-\text{Cl} \\ \quad \\ \text{H} \quad \text{H} \end{array} $	chloroethane
... [x - y] _n ...	polymer	$ \begin{array}{c} \left[\begin{array}{cc} \text{H} & \text{H} \\ & \\ -\text{C} & -\text{C}- \\ & \\ \text{H} & \text{H} \end{array} \right]_n \dots \end{array} $	polyethene
R usually represents a carbon group R' usually represents a different carbon group Q represents a halogen (fluoro-, chloro-, bromo-, iodo-)		x-y represents the monomer unit n represents a whole number	

Solutions

Solubility of Selected Ionic Compounds in Aqueous Solutions at 25°C

Ion	H ⁺ , Na ⁺ , K ⁺ , NH ₄ ⁺ , NO ₃ ⁻ , ClO ₃ ⁻ , ClO ₄ ⁻ , CH ₃ COO ⁻	F ⁻	Cl ⁻ Br ⁻ I ⁻	SO ₄ ²⁻	PO ₄ ³⁻ SO ₃ ²⁻ CO ₃ ²⁻	IO ₃ ²⁻ OOC ⁻ COO ²⁻	S ²⁻	OH ⁻
Solubility greater than or equal to 0.1 mol/L (very soluble) (aq)	most	most	most	most	H ⁺ Na ⁺ K ⁺ NH ₄ ⁺	Li ⁺ Co(IO ₃) ₂ Fe ₂ (OOC ⁻ COO) ₃	Li ⁺ Mg ²⁺ Ca ²⁺	Li ⁺ Sr ²⁺
Solubility less than 0.1 mol/L (slightly soluble) (s)	RbClO ₄ CsClO ₄ AgCH ₃ COO	Li ⁺ Mg ²⁺ Ca ²⁺ Sr ²⁺ Ba ²⁺ Fe ²⁺ Pb ²⁺	Cu ⁺ Ag ⁺ Hg ²⁺ Pb ²⁺	Ca ²⁺ Sr ²⁺ Ba ²⁺ Pb ²⁺ Ag ⁺	most	most	most	most

Note: This solubility table is only a guideline that was established using the K_{sp} values. A concentration of 0.1 mol/L corresponds to approximately 10 g/L to 30 g/L, depending on molar mass.

Stoichiometry and Solution Formulas

$$n = \frac{m}{M}$$

$$C = \frac{n}{V}$$

$$C_i V_i = C_f V_f$$

$$\frac{\text{coefficient}_r}{\text{coefficient}_g} = \frac{n_r}{n_g} \quad \text{or} \quad n_r = n_g \times \frac{\text{coefficient}_r}{\text{coefficient}_g}$$

$$(\% V/V) = \frac{V_{\text{solute}}}{V_{\text{solution}}} \times 100\%$$

$$\text{parts per million} = \frac{m_{\text{solute}}}{m_{\text{solution}}} \times 10^6 \text{ ppm}$$

n = number of moles (mol)

m = mass (g)

M = molar mass (g/mol)

C = molar concentration (mol/L)

V = volume (L)

i = initial solution

f = final solution

r = required substance

g = given substance

$\%V/V$ = percent by volume concentration

Identification of Selected Ions in 1.0 mol/L Aqueous Solutions

Ion	Symbol	Colour in Solution
chromate	$\text{CrO}_4^{2-}(\text{aq})$	yellow
chromium(III)	$\text{Cr}^{3+}(\text{aq})$	blue-green
chromium(II)	$\text{Cr}^{2+}(\text{aq})$	dark blue
cobalt(II)	$\text{Co}^{2+}(\text{aq})$	red
copper(I)	$\text{Cu}^+(\text{aq})$	blue-green
copper(II)	$\text{Cu}^{2+}(\text{aq})$	blue
dichromate	$\text{Cr}_2\text{O}_7^{2-}(\text{aq})$	orange
iron(II)	$\text{Fe}^{2+}(\text{aq})$	lime green
iron(III)	$\text{Fe}^{3+}(\text{aq})$	orange-yellow
manganese(II)	$\text{Mn}^{2+}(\text{aq})$	pale pink
nickel(II)	$\text{Ni}^{2+}(\text{aq})$	blue-green
permanganate	$\text{MnO}_4^-(\text{aq})$	deep purple

Identification of Selected Metals in Ionic Compounds

Ion	Symbol	Colour in Solution
barium	Ba	yellow-green
calcium	Ca	yellowish red
cesium	Cs	violet
copper	Cu	blue-green
lead	Pb	blue-white
lithium	Li	red
potassium	K	violet
rubidium	Rb	violet
sodium	Na	yellow
strontium	Sr	scarlet red

Acids and Bases

Rules for Naming Acids

Compound Name	Classical System Example				IUPAC System Example
	Acid Name	Formula	Compound Name	Acid Name	Acid Name
hydrogen <i>-ide</i>	<i>hydro-ic</i> acid	$\text{HCl}(\text{aq})$	hydrogen chlor <i>ide</i>	<i>hydrochloric</i> acid	aqueous hydrogen chloride
hydrogen <i>-ate</i>	<i>-ic</i> acid	$\text{H}_3\text{PO}_4(\text{aq})$	hydrogen phosph <i>ate</i>	phosphor <i>ic</i> acid	aqueous hydrogen phosphate
hydrogen <i>-ite</i>	<i>-ous</i> acid	$\text{H}_3\text{PO}_3(\text{aq})$	hydrogen phosph <i>ite</i>	phosphor <i>ous</i> acid	aqueous hydrogen phosphite

IUPAC Rules for Naming Inorganic Bases

Base Name	Example	
	Formula	Base Name
cation + anion	$\text{NaOH}(\text{aq})$	sodium hydroxide

pH Formulas

$$\text{pH} = -\log_{10}[\text{H}_3\text{O}^+(\text{aq})]$$

$$[\text{H}_3\text{O}^+(\text{aq})] = 10^{(-\text{pH})}$$

[] = concentration (mol/L)

Relative Strengths of Selected Acids and Bases for 0.10 mol/L Solution at 25°C


Acid Name	Acid Formula	Conjugate Base Formula
hydrochloric acid	HCl(aq)	Cl ⁻ (aq)
sulfuric acid	H ₂ SO ₄ (aq)	HSO ₄ ⁻ (aq)
nitric acid	HNO ₃ (aq)	NO ₃ ⁻ (aq)
hydronium ion	H ₃ O ⁺ (aq)	H ₂ O(l)
oxalic acid	HOOC ⁻ COOH(aq)	HOOC ⁻ COO ⁻ (aq)
sulfurous acid	H ₂ SO ₃ (aq)	HSO ₃ ⁻ (aq)
hydrogen sulfate ion	HSO ₄ ⁻ (aq)	SO ₄ ²⁻ (aq)
phosphoric acid	H ₃ PO ₄ (aq)	H ₂ PO ₄ ⁻ (aq)
orange IV	HOr(aq)	Or ⁻ (aq)
nitrous acid	HNO ₂ (aq)	NO ₂ ⁻ (aq)
hydrofluoric acid	HF(aq)	F ⁻ (aq)
methanoic acid	HCOOH(aq)	HCOO ⁻ (aq)
methyl orange	HMo(aq)	Mo ⁻ (aq)
benzoic acid	C ₆ H ₅ COOH(aq)	C ₆ H ₅ COO ⁻ (aq)
ethanoic (acetic) acid	CH ₃ COOH(aq)	CH ₃ COO ⁻ (aq)
carbonic acid (CO ₂ (g) + H ₂ O(l))	H ₂ CO ₃ (aq)	HCO ₃ ⁻ (aq)
bromothymol blue	HBb(aq)	Bb ⁻ (aq)
hydrosulfuric acid	H ₂ S(aq)	HS ⁻ (aq)
phenolphthalein	HPh(aq)	Ph ⁻ (aq)
boric acid	H ₃ BO ₃ (aq)	H ₂ BO ₃ ⁻ (aq)
ammonium ion	NH ₄ ⁺ (aq)	NH ₃ (aq)
hydrogen carbonate ion	HCO ₃ ⁻ (aq)	CO ₃ ²⁻ (aq)
indigo carmine	HIc(aq)	Ic ⁻ (aq)
water (55.5 mol/L)	H ₂ O(l)	OH ⁻ (aq)

Acid-Base Indicators at 25°C

Indicator	Abbreviation (acid/conjugate base)	pH Range	Colour Change as pH Increases
methyl violet	HMv(aq) / Mv ⁻ (aq)	0.0 – 1.6	yellow to blue
thymol blue	H ₂ Tb(aq) / HTb ⁻ (aq)	1.2 – 2.8	red to yellow
thymol blue	HTb ⁻ (aq) / Tb ²⁻ (aq)	8.0 – 9.6	yellow to blue
orange IV	HOr(aq) / Or ⁻ (aq)	1.4 – 2.8	red to yellow
methyl orange	HMo(aq) / Mo ⁻ (aq)	3.2 – 4.4	red to yellow
bromocresol green	HBg(aq) / Bg ⁻ (aq)	3.8 – 5.4	yellow to blue
litmus	HLt(aq) / Lt ⁻ (aq)	4.5 – 8.3	red to blue
methyl red	HMr(aq) / Mr ⁻ (aq)	4.8 – 6.0	red to yellow
chlorophenol red	HCh(aq) / Ch ⁻ (aq)	5.2 – 6.8	yellow to red
bromothymol blue	HBb(aq) / Bb ⁻ (aq)	6.0 – 7.6	yellow to blue
phenol red	HPr(aq) / Pr ⁻ (aq)	6.6 – 8.0	yellow to red
phenolphthalein	HPh(aq) / Ph ⁻ (aq)	8.2 – 10.0	colourless to pink
thymolphthalein	HTh(aq) / Th ⁻ (aq)	9.4 – 10.6	colourless to blue
alizarin yellow R	HAy(aq) / Ay ⁻ (aq)	10.1 – 12.0	yellow to red
indigo carmine	HIc(aq) / Ic ⁻ (aq)	11.4 – 13.0	blue to yellow
1,3,5–trinitrobenzene	HNb(aq) / Nb ⁻ (aq)	12.0 – 14.0	colourless to orange

Genetics

Pedigree Chart



DNA Nitrogen Bases

Nitrogen Base	Abbreviation
adenine	A
cytosine	C
guanine	G
thymine	T

Alleles

Upper case—dominant
 Lower case—recessive
 Sex linked— $X^?Y$ or $X^?X^?$

DNA Base Triplets and Their Corresponding Amino Acids

		S E C O N D B A S E					
		T	C	A	G		
F I R S T B A S E	T	TTT phenylalanine	TCT serine	TAT tyrosine	TGT cysteine	T	T H I R D
		TTC phenylalanine	TCC serine	TAC tyrosine	TGC cysteine	C	
		TTA leucine	TCA serine	TAA STOP**	TGA STOP**	A	
		TTG leucine	TCG serine	TAG STOP**	TGG tryptophan	G	
	C	CTT leucine	CCT proline	CAT histidine	CGT arginine	T	
		CTC leucine	CCC proline	CAC histidine	CGC arginine	C	
		CTA leucine	CCA proline	CAA glutamine	CGA arginine	A	
		CTG leucine	CCG proline	CAG glutamine	CGG arginine	G	
	A	ATT isoleucine	ACT threonine	AAT asparagine	AGT serine	T	
		ATC isoleucine	ACC threonine	AAC asparagine	AGC serine	C	
		ATA isoleucine	ACA threonine	AAA lysine	AGA arginine	A	
		ATG methionine or START*	ACG threonine	AAG lysine	AGG arginine	G	
	G	GTT valine	GCT alanine	GAT aspartate	GGT glycine	T	
		GTC valine	GCC alanine	GAC aspartate	GGC glycine	C	
		GTA valine	GCA alanine	GAA glutamate	GGA glycine	A	
		GTG valine	GCG alanine	GAG glutamate	GGG glycine	G	

Note: This table uses base triplets from the “complementary” ($5' \rightarrow 3'$) strand of DNA.

***Note:** ATG is an initiator base triplet but also codes for the amino acid methionine.

****Note:** TAA, TAG, and TGA are terminator base triplets.

Scoring Descriptions for Standards Setting

Standard of Excellence and Acceptable Standard for Knowledge

Score	Scoring Description
4 Standard of Excellence	The response is well organized and addresses all the major points of the question using appropriate and clear communication strategies. The description of relevant scientific, technological, and/or societal concepts is explicit . Descriptions, explanations, and/or interrelationships between the concepts provided are correct and reflect a thorough understanding of the question.
2 Acceptable Standard	The response is generally organized and addresses most of the major points of the question using adequate communication strategies. The description of relevant scientific, technological, and/or societal concepts is mentioned . Descriptions between the concepts provided are generally correct and reflect an adequate understanding of the question.

Standard of Excellence and Acceptable Standard for Skills

Score	Scoring Description
4 Standard of Excellence	The problem is thoroughly understood . An appropriate and practical design is presented. The data are accurately and completely analyzed. Accurate interpretations and conclusions are made based on an analysis of the data . The evaluation of the overall study is based on a thorough understanding of the principles of scientific inquiry.
2 Acceptable Standard	The problem is understood . The design is generally appropriate , or a practical procedure with some omissions or errors is presented. The data are adequately analyzed. Interpretations and conclusions are generally based on an analysis of the data . The evaluation of the overall study is based on an adequate understanding of the principles of scientific inquiry.

Standard of Excellence and Acceptable Standard for STS

Score	Scoring Description
4 Standard of Excellence	The design and function of the technological device are clearly explained . The interrelationships between science, technology, and society are thoroughly understood . Risks and benefits are thoroughly evaluated . Insightful and convincing arguments are used to support a decision or judgement, and a range of viewpoints is considered .
2 Acceptable Standard	The design and function of the technological device are described . The interrelationships between science, technology, and society are generally understood . Risks and benefits are listed . Logical arguments are used to support a decision or judgement, and viewpoints are considered .

References

Lide, D.R. 2005. *CRC Handbook of Chemistry and Physics*. 86th ed. Boca Raton: CRC Press.
NIST Reference on Constants, Units and Uncertainty. 2002. <http://physics.nist.gov>